CPV Treatment Flowchart

CPV-infected dog ≥5 lb, BCS ≥3/9, ≤10% dehydration, quiet/alert/responsive

Move to Isolation

- **Body weight (kg)**
- **Insert IV catheter in cephalic vein**
- **Start IV LRS fluids at 25 mL/kg/hr for 24 hr**
- **Supplement LRS fluids with KCl and 50% dextrose**
- **Cerenia, Convenia, Baytril, Simbadol SQ**
- **Keep warmed rice bags around patient**
- **Monitor temp, HR, RR, MM, CRT during fluid therapy**

Days 2 and 3
- **Body weight (kg) SID**
- **Temp, HR, RR, MM, CRT, % dehydration BID**
- **Remove IV catheter**
- **Switch to SQ LRS fluid therapy TID**
- **Cerenia, Baytril, Simbadol SQ SID**
- **Karo syrup PO TID**
- **Tumil-K PO TID**
- **a/d slurry PO TID**
- **Keep warmed rice bags around patient**
- **Monitor vomiting/diarrhea**

Alert vet if dog has:
- T>104
- T<100
- 10% loss in body weight
- Persistent dehydration >5%
- White MM, CRT>4 sec
- Rapid HR, rapid RR
- Cold legs and feet
- Altered mentation/activity
- Protracted vomiting after 2 days
- Profuse diarrhea after 3 days

Days 4 to 7
- **Body weight (kg) SID**
- **Temp, HR, RR, MM, CRT, % dehydration BID**
- **Karo syrup PO TID (if not eating)**
- **Tumil-K PO TID (if not eating)**
- **a/d slurry PO TID (if not eating)**
- **Keep warmed rice bags around patient**

Recovery
- **Alert/active**
- **Soft or normal stools**
- **Eating voluntarily**
- **Give Strongid**
- **Perform SNAP CPV test**

CPV Positive
- **Stay in Isolation**
- **No treatments**
- **Repeat CPV test daily**

CPV Negative
- **Accel shampoo bath**
- **Move out of Isolation**